

Low R_V from circumstellar dust around supernovae

Ariel Goobar

*Physics Department, Stockholm University, AlbaNova University Center, SE 106 192
Stockholm, Sweden*

ABSTRACT

The effective extinction law for supernovae surrounded by circumstellar dust is examined by Monte-Carlo simulations. Grains with light scattering properties as for interstellar dust in the Milky-Way (MW) or the Large Magellanic Clouds (LMC), but surrounding the explosion site would cause a semi-diffusive propagation of light up to the edge of the dust shell. Multiple scattering of photons predominantly attenuates photons with shorter wavelengths, thus steepening the effective extinction law as compared to the case of single scattering in the interstellar medium. Our simulations yield typical values for the total to selective extinction ratio $R_V \sim 1.5 - 2.5$, as seen in recent studies of Type Ia supernova colors, with further stiffening differential extinction toward the near-UV.

Subject headings: dust,extinction —supernovae:general

1. Introduction

The uncertainties in the brightness corrections of Type Ia supernovae (SNIa) for color excess is among the largest systematic uncertainties in the use of Type Ia supernovae to measure cosmological distances (Nordin et al. 2008). The standard interpretation of color excess being due to extinction by interstellar dust in the supernova host galaxy has recently been challenged by the empirically deduced color-brightness relation for SNIa. While studies of differential extinction of quasars shining through foreground galaxies yield values of $R_V = A_V/E(B - V)$ compatible with the average MW value (Östman et al. 2008)¹, the SNIa Hubble diagram scatter is minimized for values significantly smaller than $R_V = 3.1$ (Astier et al. 2006; Kowalski et al 2008). Furthermore, the use of SNIa for cosmology benefits significantly from the understanding the extinction law for the full optical and near-IR

¹It should be noted that low R_V values to individual QSO systems have been found in e.g. (Östman et al. 2008) and (Wang et al. 2004)

wavelength range. E.g. the wavelength dependence of extinction toward the Magellanic Clouds differ from the Milky-Way extinction law in (Cardelli et al. 1989), even for very similar values of R_V . For quasar sight-lines, Östman et al. (2008) tested both Milky-Way like extinction as well as Small Magellanic Cloud (SMC) extinction law, both giving comparable goodness of fit. A preference for SMC dust for extinction of AGNs has been suggested by Li (2007).

Recently, the detection of circumstellar (CS) matter in the local environment surrounding the Type Ia supernova SN2006X in the nearby galaxy M100 has been reported by Patat et al. (2007). A shell within a few 10^{16} cm (~ 0.01 pc) of the center of the explosion has been suggested to explain the time-variable Na I D lines in the SN spectrum.

Wang et al. (2008a) report $R_V = 1.48 \pm 0.06$ and $E(B - V) = 1.42 \pm 0.04$ mag for SN2006X and a light echo in the lightcurve was found by Wang et al. (2008b) consistent with dust illuminated at a distance of 27-170 pc from the site of the explosion. Even if the local environment around this supernova may not be very common among SNIa, similar values for the total to selective extinction ratio have been reported for several SNIa with good wavelength coverage. E.g. Krisciunas et al. (2007) found $R_V = 1.55 \pm 0.08$ for SN 1999cl; Elias-Rosa et al. (2006, 2008) report $R_V = 1.80 \pm 0.19$ and $R_V = 1.59 \pm 0.07$ for SN2003cg and SN 2002cv respectively. Furthermore, a statistical study of optical colors of a sample including 80 near-by SNIa, Nobili & Goobar (2008) found an average value of $\bar{R}_V = 1.75 \pm 0.27$ for SNIa with $E(B - V) < 0.7$, and even lower for a subsample of low-reddening SNIa.

Next, we examine the possibility that low values of R_V stem from the semi-diffusive propagation of photons in the neighborhood of the site of the supernova explosion.

2. Quasi-diffusive light propagation around the supernova

Photon propagation around a medium of scatterers can be described by a quasi random-walk picture. The reader is referred to (Chandrasekhar 1943) for a beautiful introduction to this subject. Lets consider a localized distribution of dust particles within a distance R_{CS} from the explosion site, negligibly small compared to the distance to the observer, d , i.e. $R_{SN} < R_{CS} \ll d$, where R_{SN} corresponds to the radius from where the SN radiation emerges. The trajectory of a photon will be straight until it hits a dust particle at which point the photon can either be scattered or absorbed. If the photon is scattered in a new direction, it follows a straight path until the next encounter, and so on until $r > R_{CS}$. The mean free path between interactions, λ_{eff} , is thus determined by the number density of scatterers, n ,

and their effective cross-section for scattering and absorption of light, $\sigma_{eff} = \sigma_s + \sigma_a$:

$$\lambda_{eff} = \frac{1}{n \cdot \sigma_{eff}} \quad (1)$$

For $\lambda_{eff} \gg R_{CS}$, the single scattering approximation is valid and the light beam reaching the observer is attenuated as $e^{-\tau}$, with $\tau = R_{CS}/\lambda_{eff}$. This is the case applicable for extinction by dust in the interstellar medium.

For $\lambda_{eff} \ll R_{CS}$, corresponding to a (local) high number density of scatterers, the situation is different. If the absorption probability is much lower than for scattering, photon propagation is diffusive and the average properties can be described analytically using the formulas for random-walk (Askebjerg et al. 1997). The case we are considering here is for $\lambda_{eff} \sim R_{CS}$, where the scattering cross-section exceeds the absorption cross-section, $\sigma_s > \sigma_a$, in a wavelength dependent manner. In particular, we examine the cases where the light scattering properties of dust particles in the CS matter are similar to what has been modeled for interstellar dust grains in the Milky-Way (Draine 2003) or the LMC (Weingartner & Draine 2001).

Table 1 shows the wavelength dependent albedo factor ($= \sigma_s/(\sigma_s + \sigma_a)$) and the average of the cosine of the scattering angle for interactions between light and dust particles. Also tabulated is the absorption cross-section divided by dust mass. Note that the Milky-Way parameters correspond to a dust size distribution matching $R_V = 3.1$ for dimming of stars in the Galaxy. Wang (2005) considered the impact of circumstellar dust upon the measured value of R_V , but only the extreme case where all scattered photons may reach the observer. That assumption overlooks an important aspect of the problem: while the bluer photons scatter more, they also are more likely to be absorbed. This leads to a steeper wavelength dependence of the effective extinction law, possibly explaining the unusual total to selective extinction ratios found in studies of SNIa.

3. Monte-Carlo simulation of light propagation around supernova

In order to estimate the net effect of scattering and absorption on the light reaching the outer edge of a shell of circumstellar dust around the SN site, R_{CS} , a Monte-Carlo simulation was performed. Photons with energies corresponding to the central wavelengths of the *UBVRIJHK* photometric system were generated and subsequently followed as they

propagate in the dusty medium.² A uniform distribution of scatterers within a sphere of radius R_{CS} is used in the calculations. Our treatment is rather insensitive of the physical size of R_{CS} since what governs the differences in path-lengths of photons at different wavelengths are the optical depths, $\tau_s = R_{CS}/\lambda_s$ and $\tau_a = R_{CS}/\lambda_a$. Thus, for a fixed color excess, a larger R_{CS} can be compensated by a lower number density, n , thus keeping τ_s and τ_a unchanged. Figure 1 shows the wavelength dependence of τ_a and τ_s in our calculation for MW and LMC dust types for a reddening at $r = R_{CS}$ of $E(B - V) = 0.1$. The key observation is that at short wavelengths, both τ_s and τ_a increase, while τ_s dominates. Since they scatter more, photons with short wavelength leaving the CS region to eventually reach the observer must propagate a larger path-length than photons at longer wavelengths. However, since τ_a also increases at shorter wavelengths, the relative attenuation of bluer photons is enhanced, thus generating an effective extinction law with steeper wavelength dependence, i.e. lower R_V .

SMC-type dust, as parameterized in (Weingartner & Draine 2001), has a smaller albedo in the optical to near-IR region (see Fig. 23 in Weingartner & Draine (2001)), leading to a larger value of τ_a compared to LMC or MW dust. Thus, SMC dust is not suitable to explain the anomalous SN colors with the scenario presented in this work.

For each photon starting at $R_{SN} \ll R_{CS}$, straight path-lengths between interactions (L_s, L_a) were generated from an exponential distributions $\lambda_s^{-1} \exp(-L_s/\lambda_s)$ and $\lambda_a^{-1} \exp(-L_a/\lambda_a)$, where the mean free path for scattering and absorption, $\lambda_s = (n \cdot \sigma_s)^{-1}$ and $\lambda_a = (n \cdot \sigma_a)^{-1}$ were calculated from the wavelength dependent parameters in Table 1. The scattering angle for each interaction was generated in the simulation following the Henyey-Greenstein approximation (Henyey & Greenstein 1941):

$$\frac{d\sigma}{d(\cos(\theta))} = \frac{1 - g^2}{(1 + g^2 - 2g \cos(\theta))^{\frac{3}{2}}}, \quad (2)$$

where $g = \langle \cos(\theta) \rangle$ (also listed in Table 1). The density of dust in the shell was varied to cover a wide range of reddening at $r = R_{CS}$, $0 \leq E(B - V) \leq 0.4$. The probability for photons to reach R_{CS} without absorption is calculated by repeating the ray-tracing Monte-Carlo $5 \cdot 10^5$ times for each wavelength.

²In this study we neglect the effect noted by Wang (2005) where the effective wavelength in each filter is changing with time following the color evolution of SNIa

4. Low R_V from simulations

Figure 2 shows that the attenuation of light after the circumstellar shell yields low values of $R_V = A_V/E(B-V)$, as anticipated. In particular, the simulations using LMC dust, result in an effective $R_V = 1.65$, compatible with the best fit result of (Nobili & Goobar 2008). We note, however, that the CCM extinction law (Cardelli et al. 1989) does not accurately reproduce the wavelength dependence of photons attenuation over the entire optical range, as shown in Figure 3³. The deviations are significant for the U-B color, in agreement with the observations reported in (Guy et al. 2005).

Instead, a power-law relation gives a good fit to the simulations (see Figure 4):

$$\frac{A_\lambda}{A_V} = 1 - a + a \left(\frac{\lambda}{\lambda_V} \right)^p, \quad (3)$$

where $\lambda_V = 0.55 \mu\text{m}$ is the central wavelength of the V-band filter. Since the attenuation at B-band (central wavelength λ_B) is given by $(R_V + 1)E(B - V)$, the parameters in Equation (3) are related to R_V as

$$R_V = a^{-1} \left[\left(\frac{\lambda_B}{\lambda_V} \right)^p - 1 \right]^{-1} = \frac{1}{a(0.8^p - 1)} \quad (4)$$

It is encouraging for the use of Type Ia SNe for precision cosmology that the reddening corrections may take a simple and general analytical form, if the model is confirmed. Further studies of SN colors, also including near-IR data, should be used to test this relation.

5. Summary and conclusions

Simple simulations show that circumstellar material, detected in at least one Type Ia supernova, SN2006X (Patat et al. 2007; Wang et al. 2008b), could potentially explain the empirically determined extinction law for low redshift SNIa, especially if the circumstellar material resembles LMC dust grains. Adopting the CS shell size of Patat et al. (2007), $R_{CS} \sim 10^{16}$ cm, we find that for $\tau \sim 1$, the required mass in dust around the supernova is $M_{dust} \sim 4\pi R_{CS}^2/(\sigma_a/m_{dust}) \sim 10^{-4}M_\odot$ when inserting typical values of the absorption cross-section from Table 1.

³The validity of the simulation code was tested by accurately reproducing the standard extinction laws when the diffusive process was switched off.

A simple power-law expression is found to fit very well the effective extinction law for dust in the CS environment of the supernova produced by Monte-Carlo simulations.

Depending on the thickness of the CS shell, shifts in the time of lightcurve maximum may be expected for different bands since the amount of quasi random-walk will differ. In particular, photons in redder bands will suffer less scattering and thereby less time delay. This effect should correlate with the measured reddening, $E(B - V)$. The assumption of a uniform density is not expected to be critical for the results at first order. However, a second order effect may be expected since a large scale of R_{CS} would result in a longer time for photons being “trapped” in the scattering sphere. As the the intrinsic colors of Type Ia change on a time scale of days (Nobili & Goobar 2008), time delays of photons of that time scale would affect the measured colors as a function of time.

In a forthcoming paper, potential direct observables from interaction between photons and dust will be investigated. Also, the sensitivity of the effective extinction law to the dust grain sizes and density profile in CS medium and the combination of both scattering in the circumstellar material and the interstellar medium needs to be further investigated. If the presence of CS material is indeed the source of the color-brightness relation found in SNIa, the case for restframe near-IR observations is further strengthened: the peak magnitude corrections, and their model dependence, are smaller than at optical wavelengths.

I am grateful to Vallery Stanishev for discussions triggering this work and to the anonymous referee for providing constructive comments which have greatly improved the paper. It is a pleasure to thank Christian Walck for making his random number generation package available to me. Thanks also to Eric Linder for pointing out a typo in the original manuscript.

REFERENCES

- Askebjerg, P., et al., 1997, *Applied Optics*, 36, 4168
- Astier, P., et al., 2006, *A&A*, 447, 31
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Chandrasekhar, S., 1943, *Rev. Mod. Phys.*, 15, 1
- Draine, B.T. 2003, *ApJ*, 598, 1017
- Elias-Rosa, N., et al., 2006, *MNRAS*, 369, 1880

- Elias-Rosa, N., et al., 2008, MNRAS, 384, 107
- Guy, J., Astier, P., Nobili, S., Regnault, N., & Pain, R. 2005, A&A, 443, 781
- Heney, L. C., & Greenstein, J. L., 1941, AJ, 93, 70
- Kowalski, M. et al., 2008, ApJ, in press, arXiv:0804.4142
- Krisciunas, K. et al., 2007, AJ, 133, 58
- Li, A., 2007, ASP Conf. Ser., Ho, L. C. and Wang, J.-W.(eds), Vol. 373, 561.
- Nobili, S. and Goobar, A., 2008, A&A, 487, 19
- Nordin, J., Goobar, A., & Jonsson, J. 2008, JCAP, 02, 008
- Östman, L., Goobar, A., Mörtzell, E., 2008, A&A, 485, 403
- Patat, F., et al., 2007, Science, 317, 924
- Wang, J. et al, 2004, ApJ, 609, 589
- Wang, L. 2005, ApJ, 635, L33
- Wang, X. F. et al, 2008a ApJ, 675, 626
- Wang, X. F. et al, 2008b ApJ, 677, 1060
- Weingartner, J.C., & Draine, B.T. 2001, ApJ, 548, 296




Fig. 1.— Optical depth for scattering ($\tau_s = R_{CS}/\lambda_s$) and absorption ($\tau_a = R_{CS}/\lambda_a$) as a function of wavelength from the Monte-Carlo simulation of photon propagation in a dusty circumstellar material around the SN for $E(B - V) = 0.1$ at $r = R_{CS}$. Scattering properties for average MW dust from (Draine 2003) and LMC dust (Weingartner & Draine 2001) tabulated in Table 1 were used. The bluer photons scatter more and since they also have a larger probability for absorption, fewer make it to the edge of the dusty shell, R_{CS} . As a result, a steeper wavelength dependence for attenuation is to be expected.




Fig. 2.— $R_V = A_V/E(B - V)$ vs $E(B - V)$ resulting from the Monte-Carlo simulation of photon propagation in a dusty circumstellar material around the explosion site. The “wiggles” in the curve are compatible with the statistical uncertainty of the Monte-Carlo simulation. Scattering properties for average MW dust from (Draine 2003) and LMC dust (Weingartner & Draine 2001) tabulated in Table 1 were used. The average total to selective extinction ratios found were $R_V = 2.56$ for Milky-Way type dust and $R_V = 1.65$ for dust compatible with properties in the Large Magellanic Clouds.




Fig. 3.— The extinction law, A_λ/A_V , derived from the Monte-Carlo simulation of circumstellar dust are compared with the parameterized Milky-Way extinction law (Cardelli et al. 1989). scattering properties for average MW dust from (Draine 2003) and LMC dust (Weingartner & Draine 2001) tabulated in Table 1 were used. The two cases considered yield a poor match to the standard $R_V = 3.1$ case. Even when the R_V is adjusted to fit the $A_V/E(B - V)$, differences are found in the wavelength dependence, especially at shorter wavelengths.




Fig. 4.— Empirical functional fit to the simulated data. scattering properties for average MW dust from (Draine 2003) and LMC dust (Weingartner & Draine 2001) tabulated in Table 1 were used.

Table 1. Scattering parameters for optical and near-IR photons from (Weingartner & Draine 2001; Draine 2003) corresponding to interstellar extinction in the Large Magellanic Clouds (LMC) and Milky-Way (MW) with $R_V = 3.1$.

wavelength (μm)	albedo		$g = \langle \cos(\theta) \rangle$		$\frac{\sigma_a}{m_{dust}} (\text{cm}^2/\text{g})$		filter
	MW	LMC	MW	LMC	MW	LMC	
0.36	0.6203	0.6532	0.5695	0.6072	1.595E+04	1.162E+04	U
0.44	0.6529	0.7159	0.5654	0.6153	1.191E+04	7.542E+03	B
0.55	0.6735	0.7631	0.5382	0.6059	8.551E+03	4.666E+03	V
0.65	0.6745	0.7785	0.4995	0.5815	6.722E+03	3.319E+03	R
0.80	0.6576	0.7686	0.4381	0.5289	5.170E+03	2.416E+03	I
1.22	0.5846	0.7029	0.2893	0.3892	3.224E+03	1.550E+03	J
1.63	0.5200	0.6472	0.2086	0.2911	2.351E+03	1.237E+03	H
2.19	0.4391	0.5599	0.1310	0.1792	1.670E+03	1.041E+03	K